



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

1. Proposed by Professor G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.

Three persons A, B, C , throw with three dice. They each stake \$10.00 and the one who first throws at least ten with the three dice takes the whole stake. Find the expectation of each.

Solution by the Proposer.

The chance of throwing respectively 10, 11, 12, ..., 18, with three dice is

$$\frac{27}{216}, \frac{27}{216}, \frac{25}{216}, \frac{21}{216}, \frac{15}{216}, \frac{10}{216}, \frac{6}{216}, \frac{3}{216}, \frac{1}{216}$$

The chance of throwing at least ten is equal to the sum of all these chances = $\frac{135}{216} = \frac{5}{8}$.

The chance that A will win = $\frac{5}{8}$, that he will not win = $\frac{3}{8}$. The chance that B will win = $\frac{5}{8} \cdot \frac{3}{8}$, that he will not win = $(\frac{3}{8})^2$. The chance that C will win = $\frac{5}{8} \cdot (\frac{3}{8})^2$. The chance that A will win the second throw = $\frac{5}{8} \cdot (\frac{3}{8})^3$, that B will win = $\frac{5}{8} \cdot (\frac{3}{8})^4$, that C will win = $\frac{5}{8} \cdot (\frac{3}{8})^5$.

$$\therefore A's \text{ chance} = \frac{5}{8} + \frac{5}{8} \left(\frac{3}{8}\right)^3 + \frac{5}{8} \left(\frac{3}{8}\right)^5 + \dots = \frac{5}{4},$$

$$B's \text{ chance} = \frac{5}{8} \left(\frac{3}{8}\right) + \frac{5}{8} \left(\frac{3}{8}\right)^4 + \frac{5}{8} \left(\frac{3}{8}\right)^7 + \dots = \frac{5}{4},$$

$$C's \text{ chance} = \frac{5}{8} \left(\frac{3}{8}\right)^2 + \frac{5}{8} \left(\frac{3}{8}\right)^5 + \frac{5}{8} \left(\frac{3}{8}\right)^8 + \dots = \frac{5}{7},$$

$$A's \text{ expectation} = \left(\frac{5}{4} \times 30\right) - 10 = \$9\frac{1}{4},$$

$$B's \text{ expectation} = \left(\frac{5}{4} \times 30\right) - 10 = -\$2\frac{1}{4},$$

$$C's \text{ expectation} = \left(\frac{5}{7} \times 30\right) - 10 = -\$7\frac{2}{7}.$$

Also solved by L. V. Roy, P. H. Philbrick, J. F. W. Scheffer, H. C. Whitaker, and W. H. Draughon

PROBLEMS.

4. Proposed by G. B. M. ZERR, Principal of High School, Staunton, Virginia.

Four points taken at random in each half, made by the transverse axis, of an ellipse, are joined in such a way by straight lines as to enclose an octagonal surface; find the mean area of this surface.

5. Proposed by DE VOLSON WOOD, M. A., C. E., Professor of Mechanical Engineering, Stevens Institute of Technology, Hoboken, New Jersey.

An actual case suggested the following:

An equal number of white and black balls of equal size are thrown into a rectangular box, what is the probability that there will be contiguous contact of white balls from one end of the box to the opposite end? As a special example, suppose there are 30 ball in the length of the box, 10 in the width, and 5 (or 10) layers deep.

Solutions to these problems should be received on or before May 1st.

MISCELLANEOUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

1. Proposed by G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.

To divide the arc of a cycloid into eight equal parts.